Upper Basin Storage Studies

Explanation: The actual amount of wetland storage that is available in the Devils Lake basin has been a matter of some debate in recent years. The following is a brief summary of the studies that have been conducted by various agencies or teams of agencies examining the acreage, and storage potential of wetlands in the Devils Lake basin.

1) Wetland Acreage and Storage Studies

Author: Ludden, A.P., D.L. Frink, and D.H. Johnson

Year: 1983

Title: Water storage capacity of natural wetland depressions in the Devils Lake Basin of North Dakota

Publisher: Journal of Soil and Water Conservation 38:1

Estimate of Wetlands Acreage (Drained and Undrained) in the DL Basin: Findings:

(Remaining: 412,000 acres)

Estimate of Wetland Storage in the DL Basin (Drained and Undrained) in the DL Basin:

2-Year Event: 116,000 acre-feet 100-Year Event: 435,000 acre-feet

Average Depth of Wetlands: 19.1 inches (1.59 feet) *Annotated in article*

Maximum Total Storage Remaining (Drained and Undrained): 657,000 acre-feet

Average Annual Evapotranspiration (Based on ND PPR): 29.9 inches

Limitations: Did not examine the role of evapotranspiration, climate, and antecedent moisture in storage. Did not consider storage of waterbodies larger than one 1/4 section (160 acres). Assumed uniform volume for wetlands. Limited topographical resolution. Evapotranspiration estimated based upon average for North Dakota prairie pothole region, not specific to Devils Lake. Storage of anthropogenic depressions was eliminated or reduced.

b)

Author: W.M. Schuh (NDSWC)

Year: 1996

Title: Drained and Partially Drained Wetlands in the Devils Lake Basin

Publisher: NDSWC Memorandum to Stan Hanson

Estimate of Drained Wetlands in the Devils Lake basin (207 samples in DL basin): Findings:

> Completely Drained Wetlands: 28,869 acres Partially Drained Wetlands: 40,521 acres Total Drained Wetlands (95% CI) 69,390 acres

Limitations: Did not estimate wetland acreage lost between cessation of NWI sampling and "Swampbuster."

Limited topographical resolution.

Author: Sprynczynatyk, D.A. (NDSWC)

Year: 1996

Title: Letter to Steve Blomeke (Natl. Wildlife Federation) Publisher: The North Dakota State Water Commission (Letter)

Findings:

Summary of Storage for Three Different Frequency Rain Events:

2 Year: 7.9 inches-14,912 acre-feet of storage 10 Year: 11.8 inches-22,143 acre-feet of storage 25 Year: 14.6 inches-27,340 acre-feet of storage

Limitations: Storage estimates were based upon wetland being completely dry prior to event. Limited

topographical resolution.

d)

Author: Sprynczynatyk, D.A. (NDSWC), and A.J. Sapa (USFWS)

Year: 1997

Title: Untitled NDSWC & USFWS letter

Publisher: Letter to Steve Blomeke (Natl wildlife Federation) and Roger Beaver, (DLBJWRB)

Findings: Estimate of Undrained Wetland Acres Remaining as of 1979(NWI): 252,460 acres Minimum Estimate of Altered Wetlands As of 1979(NWI)

Completely Drained: 25,490 acres Partially Drained: 34,000 acres

Assumed Wetland Loss Between 1979 and "Swampbuster" Act: 41,000 acres Total Wetlands Altered as of 1985(Personal Calculation 1): 100,490 acres

Total Wetland Area(Personal Calculation 2): 352,950 acres

Estimated Drained Wetland Storage: Based on 60,000 acres (w/out post 1979 additional 41,000 acres) drained wetland acres, utilizing Ludden(1983), Best(1978)Range of 156,000-294,000 acre-feet

Additional Storage: Available above current on upper basin lakes: 33,250 acre-feet Total Storage(Not Incorporating Wetland Loss 1979-1985): Range of 189,250-327,250 acre-feet (Personal Calc. of Min. Estimate)

Limitations: Did not extrapolate lost storage based upon additional acreage. Relied exclusively upon literature review, with no ground-truthing. Did not examine the role of evapotranspiration, climate, and antecedent moisture in storage. Limited topographical resolution.

e)

Author: Sapa, A.J. (USFWS)

Year: 1997

Title: Letter to Steve Blomeke (Natl. Wildlife Federation)

Publisher: United States Fish & Wildlife Service

Findings:

Estimate of Original Wetland Acreage in the DL basin: 400,000 acres Estimate of Drained Wetland Acreage in the DL basin: 189,000 acres Estimate of Remaining Wetland Acreage in the DL basin: 211,000 acres

Estimated Range of Storage of Drained Wetlands in the DL basin: 491,400-926,100 acre-

feet

Estimate of Maximum Storage of Remaining Wetlands in the DL basin: 657,000 acre-feet Estimate Maximum Storage of Original Wetland Acreage in the DL basin (personal calculation): 1,148,400-1,583,100 acre-feet

Estimate of Drained Hydric-Soil Acreage in the DL basin as of 1980 (NRCS source): 142,000 acres

Estimate of Required Wetland Acreage Restoration to Hold 75,000 Acres: 43,000 acres

Estimated Evaporation Rate (As from Shjeflo, 1968): 25.32 inches annually

Estimated Evapotranspiration Rate (As from Ludden et al. 1983): 29.9 inches annually

Estimated Seepage (Groundwater Loss?): 7.2 inches annually

Max Wetland Depth (Ludden, 1983, with # number changes to reflect revision): 19.1 inches

101105

Estimate of Average Original Wetland Storage/Depth (Personal Calculation: Storage of Original Wetlands/Original Wet Acres): 47.5 inches

Estimate of Average Drained Wetland Storage/Depth (Personal Calculation: Min-Max Storage of Drained Wetlands/Drained Wet Acres): 31.2 inches to 58.8 inches

Limitations: Evapotranspiration estimated based upon average for North Dakota prairie pothole region, not specific to Devils Lake. Analysis not based on specific hydrological studies.

f)

Author: Gehringer, M. and M. Hove (State Water Commission)

Year: 1998

Title: Assessment of Drainage in the Devils Lake Basin Publisher: North Dakota State Water Commission

Findings:

Number of Drained Wetlands in the Devils Lake Basin (includes Stump): 23,000 Acreage of Drained Wetlands in the Devils Lake Basin (includes Stump): 41,000 acres Volume of Drained Wetlands in the Devils Lake Basin (includes Stump): 82,000 acre-feet

Limitations: Assumed an average wetland depth of two feet. Wetland data is based upon NRCS wetlands maps of almost the entire basin, and GIS delineations of those maps. Study assumes that all drained wetlands do contribute to Devils Lake or Stump Lake.

g)

Author: Bell, A., D. Eckhardt, and M. Pucherelli (Bureau of Reclamation)

Year: 1999

Title: Pilot Project: Wetlands Inventory and Drained Wetlands Water Storage Capacity Estimation for the St.

Joe-Calio Coulee Subbasin of the Greater Devils Lake Basin, North Dakota

Publisher: U.S. Department of the Interior, Bureau of Reclamation. Technical Service Center. Denver,

Colorado. Technical Memorandum No. 8260-99-02

Findings:

Number of Existing Wetlands in St. Joe-Calio Coulee: 4,309

Acreage of Existing Wetlands in the St. Joe-Calio Coulee: 21,505 acres

Average Maximum Depth of Depressions in the St. Joe-Calio Coulee: 8.6 inches

Number of Drained Wetlands in the St. Joe-Calio Coulee: 1,898 Acreage of Drained Wetlands in the St. Joe-Calio Coulee: 4,749 acres

Storage of Drained Wetlands in the St. Joe-Calio Coulee: Estimate of 3,400 acre-feet

(95% CI)

Estimate of Drained Wetland Storage for Entire DL Basin based on St. Joe-Calio Coulee

Numbers: 53,408 acre-feet (95% CI)

Estimate of Total Wetland Storage in the Entire DL Basin based on St. Joe-Calio Coulee

Numbers: 524,000 acre-feet

Estimate of Drained Wetland Storage (Total Wetland Storage-Drained Wetland

Storage) (Personal Calculation): 470,592 acre-feet

Limitations: Estimates of basin-wide wetland storage based upon small, extensively studied data subset. Difficulty in estimating drained wetland area. Limited ground-truthing. Limited topographical resolution. Did not examine the role of evapotranspiration, climate, and antecedent moisture in storage.

h)

Author: Doeing, B.J., and S.M. Forman (West Consultants, Inc. on Behalf of USACE)

Year: 2001

Title: Final Report: Devils Lake Upper Basin storage Evaluation

Publisher: West Consultants, Inc. Prepared for United States Army Corps of Engineers
Findings: Number of Possibly Intact/Undrained Wetlands in the DL Basin: 63,458

Possibly Intact/Undrained Wetland Acreage in the DL Basin: 201,990 acres

Estimated Storage of Possibly Intact/Undrained Wetlands in the DL Basin: 481,604 acre-

feet

Number of Possible Drained Wetlands in the DL Basin: 52,210 Possibly Drained Wetland Acreage in the DL Basin: 92,429 acres

Estimated Storage of Possibly Drained Wetlands in the DL Basin: 132,729 acre-feet

Total Number of Wetlands in the DL Basin: 115,668

Total Acreage of Wetlands in the DL Basin: 294,419 acres

Total Storage of Wetlands in the DL Basin: 614,333 acre-feet

Restoration of 25% of Possibly Drained Wetlands (Option B) Acreage: 19,472 acres Restoration of 25% of Possibly Drained Wetlands (Option B) Storage: 31,431 acre-feet Restoration of 50% of Possibly Drained Wetlands (Option C) Acreage: 39,681 acres

Restoration of 50% of Possibly Drained Wetlands (Option C) Storage: 63,608 acre-feet Restoration of 75% of Possibly Drained Wetlands (Option D) Acreage: 59,872 acres Restoration of 75% of Possibly Drained Wetlands (Option D) Storage: 94,850 acre-feet Restoration of 100% of Possibly Drained Wetlands (Option D) Acreage: 79,762 acres Restoration of 100% of Possibly Drained Wetlands (Option D) Storage: 127,835 acre-feet Average Annual Runoff Reduction (10 Climate Sequences Average)(Option B): 6,836 acre-feet

Average Annual Runoff Reduction (10 Climate Sequences Average)(Option C): 12,910 acre-feet

Average Annual Runoff Reduction (10 Climate Sequences Average)(Option D): 18,403 acre-feet

Average Annual Runoff Reduction (10 Climate Sequences Average)(Option B): 23,841 acre-feet

Average Annual Runoff Reduction (Wet Scenario)(Option B): 7,959 acre-feet Average Annual Runoff Reduction (Wet Scenario)(Option C): 15,643 acre-feet Average Annual Runoff Reduction (Wet Scenario)(Option B): 23,502 acre-feet Average Annual Runoff Reduction (Wet Scenario)(Option B): 31,193 acre-feet Average Evaporation from a Depression: >20 inches annually

Limitations: Evapotranspiration was likely underestimated. Did not examine the effects of climate completely enough. No ground-truthing of estimates. Wetland restoration could be optimized to increase wetland storage for Options B-D.

i)

Author: Vining, K.C. (USGS)

Year: 2002

Title: Simulation of Streamflow and Wetland Storage, Starkweather Coulee Subbasin, North Dakota, Water

Years 1981-98

Publisher: USGS, Water-Resources Investigations Report 02-4113

Findings:

Average Wetland Depth in Starkweather Coulee: 2.21 feet

Total Maximum Wetland Area in Starkweather Coulee: 30,890 acres

Total Maximum Wetland Volume in Starkweather Coulee: 68,270 acre-feet

Average Reduction in Spillage from Wetlands with Removal of Drains (1981-1998): 49%

1997-Event Reduction in Spillage from Wetlands with Removal of Drains: 20% Average Wetland Evapotranspiration-Personal Calc. (1981-1998): Approx 20 inches annually.

Limitations: Did not model hydrological effects of frozen soil. Study was limited to hydrological factors being estimated based upon one wetland analyzed per hydrological response unit. Other hydrological modeling limitations, due to an extremely complex system.