TRENDS IN OVERALL ADULT CIGARETTE USE COMPARED TO SMOKE-FREE INDOOR AIR LAWS

	Trends in Overall Adult Cigarette Use 2000 Through 2010 ¹												Areas Covered by Smoke-Free Indoor Air Laws as of December 31, 2009 ²					
												Change			Worksites and			
State	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2000-10	Rank	Full	Restaurants	Worksites	Restaurants	None
New Hampshire	25.3	24.1	23.2	21.2	21.7	20.4	18.7	19.3	17.1	15.8	16.9	-8.4	1				X	
Nevada	29.0	26.9	26.0	25.2	23.2	23.1	22.2	21.5	22.2	22.0	21.3	-7.7	2		X			
Rhode Island	23.4	23.9	22.4	22.4	21.3	19.8	19.2	17.0	17.4	15.1	15.7	-7.7	3	Χ				
lowa	23.2	22.1	23.2	21.7	20.8	20.4	21.4	19.8	18.8	17.2	16.1	-7.1	4	Χ				
Connecticut	19.9	20.6	19.4	18.6	18.1	16.5	17.0	15.4	15.9	15.4	13.2	-6.7	5					X
South Dakota	21.9	22.3	22.6	22.7	20.3	19.8	20.3	19.8	17.5	17.5	15.4	-6.5	8		X			
New York	21.6	23.2	22.3	21.6	19.9	20.5	18.2	18.9	16.8	18.0	15.5	-6.1	10	Χ				
Missouri	27.2	25.9	26.5	27.2	24.1	23.4	23.2	24.5	25.0	23.1	21.1	-6.1	14					X
North Dakota	23.2	22.1	21.5	20.5	19.9	20.1	19.5	20.9	18.1	18.6	17.4	-5.8	18			X		
Minnesota	19.8	22.2	21.7	21.1	20.7	20.0	18.3	16.5	17.6	16.8	14.9	-4.9	35	Χ				
Alaska	25.0	26.2	29.3	26.2	24.8	24.9	24.0	22.2	21.5	20.6	20.4	-4.6	36					X
Wyoming	23.8	22.2	23.7	24.6	21.7	21.3	21.6	22.1	19.4	19.9	19.5	-4.3	37					X
Louisiana	24.1	24.6	23.9	26.5	23.5	22.6	23.4	22.6	20.5	22.1	22.1	-2.0	47		Х			
Mississippi	23.5	25.3	27.3	25.6	24.5	23.6	25.1	23.9	22.7	23.3	22.9	-0.6	48					X
Montana	18.8	21.9	21.2	20.0	20.4	19.2	18.9	19.5	18.5	16.8	18.8	0.0	49	Χ				
Oklahoma	23.3	28.7	26.6	25.1	26.1	25.1	25.1	25.8	24.7	25.5	23.7	0.4	50					X
West Virginia	26.1	28.2	28.4	27.3	26.9	26.7	25.7	26.9	26.5	25.6	26.8	0.7	51					X

¹Cigarette use is based on data from the federal Centers for Disease Control and Prevention (CDC) behavioral risk factor surveillance survey included in CDC State Tobacco Activities Tracking and Evaluation (STATE) System reports of annual cigarette use among adults. The top five states had the most significant decrease in adult cigarette use from 2000 to 2010, while the bottom five states had the least change or an increase in adult cigarette use. Other states are included based on proximity to North Dakota and other factors, including similarities in population and smoking rate reductions.

²Smoke-free indoor air laws information provided from federal CDC 2009 STATE System report. Full coverage includes worksites, restaurants, and bars.