
#### **Primary Care Access and Preventive Care**




#### CDF-AD - Screening for Depression and Follow-Up Plan Ages 18 and Older


	5	
100%	Follow-Up Plan Documented Numerator:	0
80%	Screened for Depression Denominator:	16,995
8070	Codes to Document Depression Screen Follow-Up F	Plan are NOT
60%	being used:	
0070	G8431: Screening for depression is documented as	being postive and
40%	a follow-up plan is documented <b>G8510:</b> Screening for depression is documented as	negative a
	follow-up plan is not required.	negative, a
20%		
	0.0% 0.0	%
0%	Depression Screen & Follow-Up Plan Docu	umented
F	ollow-up visit with a mental health practitioner after	discharge FFY 2020
💻 N	dedian - No National Report for this measure FFY 201	9

#### **Maternal and Perinatal Health**

CCP-AD - Contraceptive Care Postpartum Women within 3 and 60 days of delivery, Ages 21-44





#### **Care of Acute and Chronic Conditions**





Median - Ages 19-50 (39 States), Ages 51-64 (38 States), Ages 19-64 (39 States) FFY 2019

**Behavioral Health Care** 













OHD-AD - Use of Opioids at High Dosage in Persons without Cancer, Ages 18 and Older




COB-AD - Concurrent Use of Opioids and Benzodiazepines Ages 18 and Older








#### SSA-AD - Adherence to Antipsychotic Medicaitons for Individuals with Schizophrenia Ages 18 and Older







	Measure	2019	re Set	2021	2022	2023	2024
	Primary Care A	ccess a	nd Preve	ntive Ca	are		
						1000	
ccs	Cervical Cancer Screening						
CHL	Chlamydia Screening in Women						
FVA	Ages 21 to 24 Flu Vaccinations for Adults Ages 18 to 64						
CDF	Screening for Depression and Follow-Up Plan: Ages 18 and up						
BCS	Breast Cancer Screening						
АВА	Adult Body Mass Index Assessment						
	Maternal	and Pe	rinatal H	ealth	Cheffend	1.11.11	
PC01	PC-01: Elective Delivery						
PPC	Prenatal and Postpartum Care:						
ССР	Postpartum Care Contraceptive Care — Postpartum						
ccw	Women Ages 21 to 44 Contraceptive Care – All Women						
	Ages 15 to 20	2000			66.2017-20		5.0. e al
	Care of Acut	e and C	hronic Co	ondition	s		
CBP	Controlling High Blood Pressure						
HPC	Comprehensive Diabetes Care: HbA1C Poor Control (>9.0%)						
DOIO	PQI 01: Diabetes Short-Term						
PQ101	Complications Admission Rate						
PQ105	PAI 05: COPD or Asthma in Older Adults Admission Rate						
PQ108	PQI 08: Heart Failure Admission Rate		9433				
PQI15	PQI 15: Asthma in Younger Adults Admission Rate						
PCR	Plan All-Cause Readmission						
AMR	Asthma Medication Ration: Ages 19 to 64						
	151004						
HVL	HIV Viral Load Suppression						
				199. A. 179. A	Status		
-	and the second	vioral H	ealth Ca	re			
IET	Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment						
MSC	Medical Assistance with Smoking and Tobacco Use Cessation						
AMM	Antidepressant Medication Management						
FUH	Follow-Up After Hospitalization for Mental Illness: Age 18 and						
	older Diabetes Screening for People						
SSD	with Schizophrenia or Bipolar Disorder who are using						r
	Antipsychotic Medications Diabetes Care for People with						
	Serious Mental Illness: HBA1c Poor						
HPCMI	Control (>9.0%)						
OHD	Control (>9.0%) Use of Opioids at High Dosage in Persons without Cancer						
	Use of Opioids at High Dosage in						
	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for						
OHD COB OUD	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency						
OHD COB	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence						
OHD COB OUD	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency Department Visit for Alcohol and						
OHD COB OUD FUA	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence Follow-Up After Emergency Department Visit for Mental Illness Adherence to Antipsychotic Medications for Individuals with						
OHD COB OUD FUA FUM	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence Follow-Up After Emergency Department Visit for Mental Illness Adherence to Antipsychotic Medications for Individuals with Schizophrenia	perienc	e of Care				
OHD COB OUD FUA FUM	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence Follow-Up After Emergency Department Visit for Mental Illness Adherence to Antipsychotic Medications for Individuals with Schizophrenia	perience	e of Care				
OHD COB OUD FUA FUM	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence Follow-Up After Emergency Department Visit for Mental Illness Adherence to Antipsychotic Medications for Individuals with Schizophrenia Exp Consumer Assessment of Healthcare Providers and Systems (CAHPS <sup>9</sup> ) Health Plan Survey 5.0H,	perienc	e of Care				
OHD COB OUD FUA FUM SAA	Use of Opioids at High Dosage in Persons without Cancer Concurrent Use of Opioids and Benzodiazepines Use of Pharmacotherapy for Opioid Use Disorder Follow-Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence Follow-Up After Emergency Department Visit for Mental Illness Adherence to Antipsychotic Medications for Individuals with Schizophrenia Consumer Assessment of Healthcare Providers and Systems						

