



The BMJ

jblock@bmj.com

Cite this as: *BMJ* 2023;380:p382<http://dx.doi.org/10.1136/bmj.p382>

Published: 23 February 2023

BMJ INVESTIGATION

Gender dysphoria in young people is rising—and so is professional disagreement

More children and adolescents are identifying as transgender and are being offered medical treatment, especially in the US—but some providers and European authorities are urging caution because of a lack of strong evidence. **Jennifer Block** reports

Jennifer Block *investigations reporter*

Last October the American Academy of Pediatrics (AAP) gathered inside the Anaheim Convention Center in California for its annual conference. Outside, several dozen people rallied to hear speakers including Abigail Martinez, a mother whose child began hormone treatment at age 16 and died by suicide at age 19. Supporters chanted the teen's given name, Yaeli; counter protesters chanted, "Protect trans youth!" For viewers on a livestream, the feed was interrupted as the two groups fought for the camera.

The AAP conference is one of many flashpoints in the contentious debate in the United States over if, when, and how children and adolescents with gender dysphoria should be medically or surgically treated. US medical professional groups are aligned in support of "gender affirming care" for gender dysphoria, which may include gonadotrophin releasing hormone analogues (GnRHa) to suppress puberty; oestrogen or testosterone to promote secondary sex characteristics; and surgical removal or augmentation of breasts, genitals, or other physical features. At the same time, however, several European countries have issued guidance to limit medical intervention in minors, prioritising psychological care.

The discourse is polarised in the US. Conservative politicians, pundits, and social media influencers accuse providers of pushing "gender ideology" and even "child abuse," lobbying for laws banning medical transition for minors. Progressives argue that denying access to care is a transphobic violation of human rights. There's little dispute within the medical community that children in distress need care, but concerns about the rapid widespread adoption of interventions and calls for rigorous scientific review are coming from across the ideological spectrum.¹

The surge in treatment of minors

More adolescents with no history of gender dysphoria—predominantly birth registered females²—are presenting at gender clinics. A recent analysis of insurance claims by Komodo Health found that nearly 18 000 US minors began taking puberty blockers or hormones from 2017 to 2021, the number rising each year.^{3,4} Surveys aiming to measure prevalence have found that about 2% of high school aged teens identify as "transgender."⁵ These young people are also more likely than their cisgender peers

to have concurrent mental health and neurodiverse conditions including depression, anxiety, attention deficit disorders, and autism.⁶ In the US, although Medicaid coverage varies by state and by treatment, the Biden administration has warned states that not covering care is in violation of federal law prohibiting discrimination.⁷ Meanwhile, the number of private clinics that focus on providing hormones and surgeries has grown from just a few a decade ago to more than 100 today.⁴

As the number of young people receiving medical transition treatments rises, so have the voices of those who call themselves "detransitioners" or "retransitioners," some of whom claim that early treatment caused preventable harm.⁸ Large scale, long term research is lacking,⁹ and researchers disagree about how to measure the phenomenon, but two recent studies suggest that as many as 20-30% of patients may discontinue hormone treatment within a few years.^{10,11} The World Professional Association for Transgender Health (WPATH) asserts that detransition is "rare."¹²

Chloe Cole, now aged 18, had a double mastectomy at age 15 and spoke at the AAP rally. "Many of us were young teenagers when we decided, on the direction of medical experts, to pursue irreversible hormone treatments and surgeries," she read from her tablet at the rally, which had by this time moved indoors to avoid confrontation. "This is not informed consent but a decision forced under extreme duress."

Scott Hadland, chief of adolescent medicine at Massachusetts General Hospital and Harvard Medical School, dismissed the "handful of cruel protesters" outside the AAP meeting in a tweet that morning. He wrote, "Inside 10 000 pediatricians stand in solidarity for trans & gender diverse kids & their families to receive evidence-based, lifesaving, individualized care."¹³

Same evidence, divergent recommendations

Three organisations have had a major role in shaping the US's approach to gender dysphoria care: WPATH, the AAP, and the Endocrine Society (see box). On 15 September 2022 WPATH published the eighth edition of its Standards of Care for the Health of Transgender and Gender Diverse People, with new chapters on children and adolescents and no minimum age requirements for hormonal and surgical treatments.^{2,12} GnRHa treatment, says WPATH, can

be initiated to arrest puberty at its earliest stage, known as Tanner stage 2.

The Endocrine Society also supports hormonal and surgical intervention in adolescents who meet criteria in clinical practice guidelines published in 2009 and updated in 2017.¹⁴ And the AAP's 2018 policy statement, *Ensuring Comprehensive Care and Support for Transgender and Gender-Diverse Children and Adolescents*, says that "various interventions may be considered to better align" a young person's "gender expression with their underlying identity."¹⁵ Among the components of "gender affirmation" the AAP names social transition, puberty blockers, sex hormones, and surgeries. Other prominent professional organisations, such as the American Medical Association, have issued policy statements in opposition to legislation that would curtail access to medical treatment for minors.¹⁶⁻¹⁹

These documents are often cited to suggest that medical treatment is both uncontroversial and backed by rigorous science. "All of those medical societies find such care to be evidence-based and medically necessary," stated a recent article on transgender healthcare for children published in *Scientific American*.²⁰ "Transition related healthcare is not controversial in the medical field," wrote Gillian Branstetter, a frequent spokesperson on transgender issues currently with the American Civil Liberties Union, in a 2019 guide for reporters.²¹ Two physicians and an attorney from Yale recently opined in the *Los Angeles Times* that "gender-affirming care is standard medical care, supported by major medical organizations . . . Years of study and scientific scrutiny have established safe, evidence-based guidelines for delivery of lifesaving, gender-affirming care."²² Rachel Levine, the US assistant secretary for health, told National Public Radio last year regarding such treatment, "There is no argument among medical professionals."²³

Internationally, however, governing bodies have come to different conclusions regarding the safety and efficacy of medically treating gender dysphoria. Sweden's National Board of Health and Welfare, which sets guidelines for care, determined last year that the risks of puberty blockers and treatment with hormones "currently outweigh the possible benefits" for minors.²⁴ Finland's Council for Choices in Health Care, a monitoring agency for the country's public health services, issued similar guidelines, calling for psychosocial support as the first line treatment.²⁵ (Both countries restrict surgery to adults.)

Medical societies in France, Australia, and New Zealand have also leant away from early medicalisation.²⁶⁻²⁷ And NHS England, which is in the midst of an independent review of gender identity services, recently said that there was "scarce and inconclusive evidence to support clinical decision making"²⁸ for minors with gender dysphoria²⁹ and that for most who present before puberty it will be a "transient phase," requiring clinicians to focus on psychological support and to be "mindful" even of the risks of social transition.³⁰

Box: The origins of paediatric gender medicine in the United States

The World Professional Association for Transgender Health (WPATH) began as a US based advocacy group and issued the first edition of the Standards of Care in 1979, when it was serving a small population of mostly adult male-to-female transsexuals. "WPATH became the standard because there was nobody else doing it," says Erica Anderson, a California based clinical psychologist and former WPATH board member. The professional US organisations that lined up in support "looked heavily to WPATH and the Endocrine Society for their guidance," she told *The BMJ*.

The Endocrine Society's guidance for adolescents grew out of clinicians' research in the Netherlands in the late 1990s and early 2000s. Peggy Cohen-Kettenis, a Utrecht gender clinic psychologist, collaborated with endocrinologists in Amsterdam, one of whom had experience of prescribing gonadotrophin releasing hormone analogues, relatively new at the time. Back then, gender dysphoric teens had to wait until the age of majority for sex hormones, but the team proposed that earlier intervention could benefit carefully selected minors.⁴⁰

The clinic treated one natal female patient with triptorelin, published a case study and feasibility proposal, and began treating a small number of children at the turn of the millennium. The Dutch Protocol was published in 2006, referring to 54 children whose puberty was being suppressed and reporting preliminary results on the first 21.⁴¹ The researchers received funding from Ferring Pharmaceuticals, the manufacturer of triptorelin.

In 2007 the endocrinologist Norman Spack began using the protocol at Boston Children's Hospital and joined Cohen-Kettenis and her Dutch colleagues in writing the Endocrine Society's first clinical practice guideline.⁴² When that was published in 2009, puberty had been suppressed in just over 100 gender dysphoric young people.⁴⁰

American Academy of Pediatrics (AAP) committee members began discussing the need for a statement in 2014, four years before publication, says Jason Rafferty, assistant professor of paediatrics and psychiatry at Brown University, Rhode Island, and the statement's lead author. "The AAP recognised that it had a responsibility to provide some clinical guidance, but more importantly to come out with a statement that said we need research, we need to integrate the principles of gender affirmative care into medical education and into child health," he says. "What our policy statement is not meant to be is a protocol or guidelines in and of themselves."

"Don't call them evidence based"

"The brief history of guidelines is that, going back more than 30 years ago, experts would write articles and so on about what people should do. But formal guidelines as we think of them now were seldom or non-existent," says Gordon Guyatt, distinguished professor in the Department of Health Research Methods, Evidence, and Impact at McMaster University, Ontario.

That led to the movement towards developing criteria for what makes a "trustworthy guideline," of which Guyatt was a part.³¹ One pillar of this, he told *The BMJ*, is that they "are based on systematic review of the relevant evidence," for which there are also now standards, as opposed to a traditional narrative literature review in which "a bunch of experts write whatever they felt like using no particular standards and no particular structure."

Mark Helfand, professor of medical informatics and clinical epidemiology at Oregon Health and Science University, says, "An evidence based recommendation requires two steps." First, "an unbiased, thorough, critical systematic review of all the relevant evidence." Second, "some commitment to link the strength of the recommendations to the quality of the evidence."

The Endocrine Society commissioned two systematic reviews for its clinical practice guideline, *Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons*: one on the effects of sex steroids on lipids and cardiovascular outcomes, the other on their effects on bone health.³²⁻³³ To indicate the quality of evidence underpinning its various guidelines, the Endocrine Society employed the GRADE system (grading of recommendations assessment, development, and evaluation) and judged the quality of evidence for all recommendations on adolescents as "low" or "very low."

Guyatt, who co-developed GRADE, found "serious problems" with the Endocrine Society guidelines, noting that the systematic reviews

didn't look at the effect of the interventions on gender dysphoria itself, arguably "the most important outcome." He also noted that the Endocrine Society had at times paired strong recommendations—phrased as "we recommend"—with weak evidence. In the adolescent section, the weaker phrasing "we suggest" is used for pubertal hormone suppression when children "first exhibit physical changes of puberty"; however, the stronger phrasing is used to "recommend" GnRHa treatment.

"GRADE discourages strong recommendations with low or very low quality evidence except under very specific circumstances," Guyatt told *The BMJ*. Those exceptions are "very few and far between," and when used in guidance, their rationale should be made explicit, Guyatt said. In an emailed response, the Endocrine Society referenced the GRADE system's five exceptions, but did not specify which it was applying.

Helfand examined the recently updated WPATH Standards of Care and noted that it "incorporated elements of an evidence based guideline." For one, WPATH commissioned a team at Johns Hopkins University in Maryland to conduct systematic reviews.^{34 35} However, WPATH's recommendations lack a grading system to indicate the quality of the evidence—one of several deficiencies. Both Guyatt and Helfand noted that a trustworthy guideline would be transparent about all commissioned systematic reviews: how many were done and what the results were. But Helfand remarked that neither was made clear in the WPATH guidelines and also noted several instances in which the strength of evidence presented to justify a recommendation was "at odds with what their own systematic reviewers found."

For example, one of the commissioned systematic reviews found that the strength of evidence for the conclusions that hormonal treatment "may improve" quality of life, depression, and anxiety among transgender people was "low," and it emphasised the need for more research, "especially among adolescents."³⁵ The reviewers also concluded that "it was impossible to draw conclusions about the effects of hormone therapy" on death by suicide.

Despite this, WPATH recommends that young people have access to treatments after comprehensive assessment, stating that the "emerging evidence base indicates a general improvement in the lives of transgender adolescents."¹² And more globally, WPATH asserts, "There is strong evidence demonstrating the benefits in quality of life and well-being of gender-affirming treatments, including endocrine and surgical procedures," procedures that "are based on decades of clinical experience and research; therefore, they are not considered experimental, cosmetic, or for the mere convenience of a patient. They are safe and effective at reducing gender incongruence and gender dysphoria."¹²

Those two statements are each followed by more than 20 references, among them the commissioned systematic review. This stood out to Helfand as obscuring which conclusions were based on evidence versus opinion. He says, "It's a very strange thing to feel that they had to cite some of the studies that would have been in the systematic review or purposefully weren't included in the review, because that's what the review is for."

For minors, WPATH contends that the evidence is so limited that "a systematic review regarding outcomes of treatment in adolescents is not possible." But Guyatt counters that "systematic reviews are always possible," even if few or no studies meet the eligibility criteria. If an entity has made a recommendation without one, he says, "they'd be violating standards of trustworthy guidelines." Jason Rafferty, assistant professor of paediatrics and psychiatry at Brown University, Rhode Island, and lead author of the AAP

statement, remarks that the AAP's process "doesn't quite fit the definition of systematic review, but it is very comprehensive."

Sweden conducted systematic reviews in 2015 and 2022 and found the evidence on hormonal treatment in adolescents "insufficient and inconclusive."²⁴ Its new guidelines note the importance of factoring the possibility that young people will detransition, in which case "gender confirming treatment thus may lead to a deteriorating of health and quality of life (i.e., harm)."

Cochrane, an international organisation that has built its reputation on delivering independent evidence reviews, has yet to publish a systematic review of gender treatments in minors. But *The BMJ* has learnt that in 2020 Cochrane accepted a proposal to review puberty blockers and that it worked with a team of researchers through 2021 in developing a protocol, but it ultimately rejected it after peer review. A spokesperson for Cochrane told *The BMJ* that its editors have to consider whether a review "would add value to the existing evidence base," highlighting the work of the UK's National Institute for Health and Care Excellence, which looked at puberty blockers and hormones for adolescents in 2021. "That review found the evidence to be inconclusive, and there have been no significant primary studies published since."

In 2022 the state of Florida's Agency for Health Care Administration commissioned an overview of systematic reviews looking at outcomes "important to patients" with gender dysphoria, including mental health, quality of life, and complications. Two health research methodologists at McMaster University carried out the work, analysing 61 systematic reviews and concluding that "there is great uncertainty about the effects of puberty blockers, cross-sex hormones, and surgeries in young people." The body of evidence, they said, was "not sufficient" to support treatment decisions.

Calling a treatment recommendation "evidence based" should mean that a treatment has not just been systematically studied, says Helfand, but that there was also a finding of high quality evidence supporting its use. Weak evidence "doesn't just mean something esoteric about study design, it means there's uncertainty about whether the long term benefits outweigh the harms," Helfand adds.

"Evidence itself never tells you what to do," says Guyatt. That's why guidelines must make explicit the values and preferences that underlie the recommendation.

The Endocrine Society acknowledges in its recommendations on early puberty suppression that it is placing "a high value on avoiding an unsatisfactory physical outcome when secondary sex characteristics have become manifest and irreversible, a higher value on psychological well-being, and a lower value on avoiding potential harm."¹⁴

WPATH acknowledges that while its latest guidelines are "based upon a more rigorous and methodological evidence-based approach than previous versions," the evidence "is not only based on the published literature (direct as well as background evidence) but also on consensus-based expert opinion." In the absence of high quality evidence and the presence of a patient population in need—who are willing to take on more personal risk—consensus based guidelines are not unwarranted, says Helfand. "But don't call them evidence based."

An evidence base under construction

In 2015 the US National Institutes of Health awarded a \$5.7m (£4.7m; €5.3m) grant to study "the impact of early medical treatment in transgender youth."³⁶ The abstract submitted by applicants said that the study was "the first in the US to evaluate longitudinal

outcomes of medical treatment for transgender youth and will provide essential evidence-based data on the physiological and psychosocial effects and safety” of current treatments. Researchers are following two groups, one of participants who began receiving GnRHa in early puberty and another group who began cross sex hormone treatment in adolescence. The study doesn’t include a concurrent no-treatment control group.

Robert Garofalo, chief of adolescent medicine at the Lurie Children’s Hospital in Chicago and one of four principal investigators, told a podcast interviewer in May 2022 that the evidence base remained “a challenge . . . it is a discipline where the evidence base is now being assembled” and that “it’s truly lagging behind [clinical practice], I think, in some ways.” That care, he explained, was “being done safely. But only now, I think, are we really beginning to do the type of research where we’re looking at short, medium, and long term outcomes of the care that we are providing in a way that I think hopefully will be either reassuring to institutions and families and patients or also will shed a light on things that we can be doing better.”³⁷

While Garofalo was doing the research he served as “contributor” on the AAP’s widely cited 2018 policy statement, which recommends that children and adolescents “have access to comprehensive, gender-affirming, and developmentally appropriate health care,” including puberty blockers, sex hormones, and, on a case-by-case basis, surgeries.¹⁵

Garofalo said in the May interview, “There is universal support for gender affirming care from every mainstream US based medical society that I can think of: the AMA, the APA, the AAP. I mean, these organisations never agree with one another.” Garofalo declined an interview and did not respond to *The BMJ*’s requests for comment.

The rush to affirm

Sarah Palmer, a paediatrician in private practice in Indiana, is one of five coauthors of a 2022 resolution submitted to the AAP’s leadership conference asking that it revisit the policy after “a rigorous systematic review of available evidence regarding the safety, efficacy, and risks of childhood social transition, puberty blockers, cross sex hormones and surgery.” In practice, Palmer told *The BMJ*, clinicians define “gender affirming” care so broadly that “it’s been taken by many people to mean go ahead and do anything that affirms. One of the main things I’ve seen it used for is masculinising chest surgery, also known as mastectomy in teenage patients.” The AAP has told *The BMJ* that all policy statements are reviewed after five years and so a “revision is under way,” based on its experts’ own “robust evidence review.”

Palmer says, “I’ve seen a quick evolution, from kids with a very rare case of gender dysphoria who were treated with a long course of counselling and exploration before hormones were started,” to treatment progressing “very quickly—even at the first visit to gender clinic—and there’s no psychologist involved anymore.”

Laura Edwards-Leeper, a clinical psychologist who worked with the endocrinologist Norman Spack in Boston and coauthored the WPATH guidelines for adolescents, has observed a similar trend. “More providers do not value the mental health component,” she says, so in some clinics families come in and their child is “pretty much fast tracked to medical intervention.” In a study of teens at Seattle Children’s Hospital’s gender clinic, two thirds were taking hormones within 12 months of the initial visit.³⁸

The British paediatrician Hilary Cass, in her interim report of a UK review into services for young people with gender identity issues, noted that some NHS staff reported feeling “under pressure to adopt

an unquestioning affirmative approach and that this is at odds with the standard process of clinical assessment and diagnosis that they have been trained to undertake in all other clinical encounters.”

Eli Coleman, lead author of WPATH’s Standards of Care and former director of the Institute for Sexual and Gender Health at the University of Minnesota, told *The BMJ* that the new guidelines emphasised “careful assessment prior to any of these interventions” by clinicians who have appropriate training and competence to assure that minors have “the emotional and cognitive maturity to understand the risks and benefits.” He adds, “What we know and what we don’t know has to be explained to youth and their parents or caregivers in a balanced way which really details that this is the evidence that we have, that we obviously would like to have more evidence, and that this is a risk-benefit scenario that you have to consider.”

Joshua Safer, director of the Center for Transgender Medicine and Surgery at Mount Sinai Hospital in New York and coauthor of the Endocrine Society guidelines, told *The BMJ* that assessment is standard practice at the programme he leads. “We start with a mental health evaluation for anybody under the age of 18,” he says. “There’s a lot of talking going on—that’s a substantial element of things.” Safer has heard stories of adolescents leaving a first or second appointment with a prescription in hand but says that these are overblown. “We really do screen these kids pretty well, and the overwhelming majority of kids who get into these programmes do go on to other interventions,” he says.

Without an objective diagnostic test, however, others remain concerned. The demand for services has led to a “perfunctory informed consent process,” wrote two clinicians and a researcher in a recent issue of the *Journal of Sex and Marital Therapy*,³⁹ in spite of two key uncertainties: the long term impacts of treatment and whether a young person will persist in their gender identity. And the widespread impression of medical consensus doesn’t help. “Unfortunately, gender specialists are frequently unfamiliar with, or discount the significance of, the research in support of these two concepts,” they wrote. “As a result, the informed consent process rarely adequately discloses this information to patients and their families.”

For Guyatt, claims of certainty represent both the success and failure of the evidence based medicine movement. “Everybody now has to claim to be evidence based” in order to be taken seriously, he says—that’s the success. But people “don’t particularly adhere to the standard of what is evidence based medicine—that’s the failure.” When there’s been a rigorous systematic review of the evidence and the bottom line is that “we don’t know,” he says, then “anybody who then claims they do know is not being evidence based.”

This feature has been funded by the BMJ Investigations Unit. For details see bmj.com/investigations.

Competing interests: I have read and understood BMJ policy on declaration of interests and have no relevant interests to declare.

Provenance: commissioned; externally peer reviewed.

- 1 Parker K, Horowitz JM, Brown A. Americans’ complex views on gender identity and transgender issues. Pew Research Center’s Social & Demographic Trends Project. 2022. <https://www.pewresearch.org/social-trends/2022/06/28/americans-complex-views-on-gender-identity-and-transgender-issues/>
- 2 Block J. US transgender health guidelines leave age of treatment initiation open to clinical judgment. *BMJ* 2022;378. doi: 10.1136/bmj.n2303 pmid: 36167353
- 3 Respaut R, Terhune C. Number of transgender children seeking treatment surges in US. *Reuters* 2022 Oct 6. <https://www.reuters.com/investigates/special-report/usa-transyouth-data/>
- 4 Terhune C, Respaut R, Conlin M. As children line up at gender clinics, families confront many unknowns. *Reuters* 2022 Oct 6. <https://www.reuters.com/investigates/special-report/usa-trans-youth-care/>

- 5 Johns MM, Lowry R, Andrzejewski J, et al. Transgender identity and experiences of violence victimization, substance use, suicide risk, and sexual risk behaviors among high school students—19 States and large urban school districts, 2017. *MMWR Morb Mortal Wkly Rep* 2019;68:71. doi: 10.15585/mmwr.mm6803a3 pmid: 30677012
- 6 Becerra-Culqui TA, Liu Y, Nash R, et al. Mental health of transgender and gender nonconforming youth compared with their peers. *Pediatrics* 2018;141:e20173845. doi: 10.1542/peds.2017-3845 pmid: 29661941
- 7 Gomez I, Ranji U, Salganicoff A, et al. Update on Medicaid coverage of gender-affirming health services. KFF. 2022 <https://www.kff.org/womens-health-policy/issue-brief/update-on-medicaid-coverage-of-gender-affirming-health-services/>
- 8 Littman L. Individuals treated for gender dysphoria with medical and/or surgical transition who subsequently detransitioned: a survey of 100 detransitioners. *Arch Sex Behav* 2021;50:69. doi: 10.1007/s10508-021-02163-w pmid: 34665380
- 9 Respaut R, Terhune C, Conlin M. Why detransitioners are crucial to the science of gender care. *Reuters* 2022 Dec 22. <https://www.reuters.com/investigates/special-report/usa-transyouth-outcomes/>
- 10 Boyd I, Hackett T, Bewley S. Care of Transgender Patients: A General Practice Quality Improvement Approach. *Healthcare (Basel)* 2022;10. doi: 10.3390/healthcare10010121. pmid: 35052285
- 11 Roberts CM, Klein DA, Adirim TA, Schvey NA, Hisle-Gorman E. Continuation of Gender-affirming Hormones Among Transgender Adolescents and Adults. *J Clin Endocrinol Metab* 2022;107:43. doi: 10.1210/clinem/dgac251. pmid: 35452119
- 12 Coleman E, Radix AE, Bouman WP, et al. Standards of care for the health of transgender and gender diverse people, version 8. *Int J Transgend Health* 2022;23(Suppl 1):259. doi: 10.1080/26895269.2022.2100644 pmid: 36238954
- 13 Hadland S. A handful of cruel protesters outside the #AAP2022 meeting, but inside 10 000 pediatricians stand in solidarity for trans & gender-diverse kids & their families to receive evidence based, lifesaving, individualized care between patients, parents & their doctor. @AmerAcadPeds: pic.twitter.com/b2K2jdRnX9. Twitter. 2022. <https://twitter.com/DrScottHadland/status/1578815082590400512>
- 14 Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine treatment of gender-dysphoric/gender-incongruent persons: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 2017;102:903. doi: 10.1210/jc.2017-01658 pmid: 28945902
- 15 Rafferty J Committee on Psychosocial Aspects of Child and Family Health Committee on Adolescence Section on Lesbian, Gay, Bisexual, and Transgender Health and Wellness. Ensuring comprehensive care and support for transgender and gender-diverse children and adolescents. *Pediatrics* 2018;142:e20182162. doi: 10.1542/peds.2018-2162 pmid: 30224363
- 16 American Academy of Child and Adolescent Psychiatry. AACAP statement responding to efforts to ban evidence-based care for transgender and gender diverse. https://www.aacap.org/AACAP/Latest_News/AACAP_Statement_Responding_to_Efforts-to_ban_Evidence-Based_Care_for_Transgender_and_Gender_Diverse.aspx
- 17 American Medical Association. March 26, 2021: State advocacy update. 2021. <https://www.ama-assn.org/health-care-advocacy/advocacy-update/march-26-2021-state-advocacy-update>
- 18 American Psychological Association. Resolution on supporting sexual/gender diverse children and adolescents in schools. 2020. <https://www.apa.org/pi/lgbt/resources/policy/gender-diverse-children>
- 19 American Psychiatric Association. Position statement on treatment of transgender (trans) and gender diverse youth. 2020. <https://www.psychiatry.org/File%20Library/About-APA/Organization-Documents-Policies/Policies/Position-Transgender-Gender-Diverse-Youth.pdf>
- 20 Boerner H. What the science on gender-affirming care for transgender kids really shows. *Sci Am* 2022 (published online 12 May). <https://www.scientificamerican.com/article/what-the-science-on-gender-affirming-care-for-transgender-kids-really-shows/>
- 21 Branstetter G. Transgender youth & health care: a guide for reporters. *Medium* 2019. <https://gillbranstetter.medium.com/transgender-youth-health-care-a-guide-for-reporters-820f8fbaff21>
- 22 Olezeski C, McNamara M, Alstott A. Op-ed: Denying trans youth gender-affirming care is an affront to science and medical ethics. *Los Angeles Times* 2022. <https://www.latimes.com/opinion/story/2022-06-13/trans-youth-healthcare-state-bans>
- 23 Simmons-Duffin S. Rachel Levine calls state anti-LGBTQ bills disturbing and dangerous to trans youth. *NPR* 2022 Apr 29. <https://www.npr.org/sections/health-shots/2022/04/29/1095227346/rachel-levine-calls-state-anti-lgbtq-bills-disturbing-and-dangerous-to-trans-you>
- 24 Socialstyrelsen: National Board of Health and Welfare. Care of children and adolescents with gender dysphoria. Report 2022-3-7799. 2022. <https://www.socialstyrelsen.se/globalassets/sharepoint-dokument/artikelkatalog/kunskapsstod/2022-3-7799.pdf>
- 25 Palveluvalikoima (Council for Choices in Health Care in Finland). Medical treatment methods for gender dysphoria in non-binary adults—recommendation. Jun 2020. https://palveluvalikoima.fi/documents/1237350/22895623/Summary_non-binary_en.pdf/8e5f9035-6c98-40d9-6acd-7459516d6f92/Summary_non-binary_en.pdf?i=1592318035000
- 26 Académie Nationale de Médecine. La médecine face à la transidentité de genre chez les enfants et les adolescents [Medicine and gender transidentity in children and adolescents. 25 Feb 2022. <https://www.academie-medecine.fr/la-medecine-face-a-la-transidentite-de-genre-chez-les-enfants-et-les-adolescents/?lang=en> (In French and English)
- 27 Royal Australian and New Zealand College of Psychiatrists. Recognising and addressing the mental health needs of people experiencing gender dysphoria/gender incongruence. 2021. <https://www.ranzcp.org/news-policy/policy-and-advocacy/position-statements/gender-dysphoria>
- 28 NHS England. Implementing advice from the Cass review. <https://www.england.nhs.uk/commissioning/spec-services/npc-crg/gender-dysphoria-clinical-programme/implementing-advice-from-the-cass-review/>
- 29 Cass Review. NICE evidence reviews. 2021. <https://cass.independent-review.uk/nice-evidence-reviews/>
- 30 NHS England. Interim service specification: specialist service for children and young people with gender dysphoria (phase 1 providers). Oct 2022. https://www.engage.england.nhs.uk/specialised-commissioning/gender-dysphoria-services/user_uploads/b1937-ii-specialist-service-for-children-and-young-people-with-gender-dysphoria-1.pdf
- 31 Institute of Medicine Board on Health Care Services Committee on Standards for Developing Trustworthy Clinical Practice Guidelines. *Clinical practice guidelines we can trust*. National Academies Press, 2011. https://play.google.com/store/books/details?id=b_RTRs8SE0YC
- 32 Maraka S, Singh Ospina N, Rodriguez-Gutierrez R, et al. Sex steroids and cardiovascular outcomes in transgender individuals: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2017;102:23. doi: 10.1210/jc.2017-01643 pmid: 28945852
- 33 Singh-Ospina N, Maraka S, Rodriguez-Gutierrez R, et al. Effect of sex steroids on the bone health of transgender individuals: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2017;102:13. doi: 10.1210/jc.2017-01642 pmid: 28945851
- 34 Wilson LM, Baker KE, Sharma R, Dukhanin V, McArthur K, Robinson KA. Effects of antiandrogens on prolactin levels among transgender women on estrogen therapy: A systematic review. *Int J Transgend Health* 2020;21:402. doi: 10.1080/15532739.2020.1819505 pmid: 34993517
- 35 Baker KE, Wilson LM, Sharma R, Dukhanin V, McArthur K, Robinson KA. Hormone therapy, mental health, and quality of life among transgender people: a systematic review. *J Endocr Soc* 2021;5:bvab011.
- 36 National Institutes of Health. The impact of early medical treatment in transgender youth. NIH Reporter. <https://reporter.nih.gov/project-details/8965408>
- 37 Northwestern University Feinberg School of Medicine. Evidence-based gender-affirming care for young adults with Robert Garofalo, MD, MPH. 20 May 2022. <https://www.feinberg.northwestern.edu/research/news/podcast/2022/evidence-based-gender-affirming%20care-for-young-adults-with-robert-garofalo-md-mp-h.html>
- 38 Tordoff DM, Wanta JW, Collin A, Stepany C, Inwards-Breland DJ, Ahrens K. Mental health outcomes in transgender and nonbinary youths receiving gender-affirming care. *JAMA Netw Open* 2022;5:e220978. doi: 10.1001/jamanetworkopen.2022.0978 pmid: 35212746
- 39 Levine SB, Abbruzzese E, Mason JW. Reconsidering informed consent for trans-identified children, adolescents, and young adults. *J Sex Marital Ther* 2022;48:27. doi: 10.1080/0092623X.2022.2046221 pmid: 35300570
- 40 Biggs M. The Dutch Protocol for Juvenile Transsexuals: Origins and Evidence. *J Sex Marital Ther* 2022:21. doi: 10.1080/0092623X.2022.2121238 pmid: 36120756
- 41 Delemarre-van de Waal HA, Cohen-Kettenis PT. Clinical management of gender identity disorder in adolescents: a protocol on psychological and paediatric endocrinology aspects. *Eur J Endocrinol* 2006;155(Suppl 1):7. https://academic.oup.com/aje/article/155/Supplement_1/S131/6695708 doi: 10.1530/eje.1.02231.
- 42 Hembree WC, Cohen-Kettenis P, Delemarre-van de Waal HA, et al. Endocrine Society. Endocrine treatment of transsexual persons: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2009;94:54. doi: 10.1210/jc.2009-0345 pmid: 19509099