

North Dakota Violent Death Reporting System Suicide Data


Data Notes: Finalized Data for 2019-2022
*Preliminary Data for 2023 and 2024

Numbers represent both ND residents and Out-of-State residents where initial injury occurred in ND


Data Source: Health Statistics and Performance

Last Updated: January 14, 2025


Suicides



Suicides by Age Group




Suicide by Sex




6.34%
Hispanic Ethnicity


Suicide by Marital Status




Suicides by Race




North Dakota Suicides




Suicides by Race





Suicides by Age Group



Suicide by Sex





Race

Select all

American Indian

Other

White

Sex

Select all

Female

Male

Age Group

Select all

10-19

20-29

30-39

40-49


50-59

60-69

70+


Suicide Location Where Injury Occurred

Year 2020 2021 2022 2023 2024



Suicides by Place of Death

Year 2020 2021 2022 2023 2024



Race



Select all American Indian Other White

Sex



☐ Select all
☐ Female
☐ Male

Age Group


Select all 10-19 20-29 30-39 40-49 50-59 60-69 70+



Suicide Method Type



Firearm Type Used



Race

Select all

American Indian

Other

White

Sex

Select all

Female

Male

Age Group

Select all

10-19

20-29


30-39

40-49

50-59

60-69

70+



0.02%: This is the lowest level of intoxication with some measurable impact on the brain and body.

0.05%: At this level behavior will become exaggerated with impaired judgement and reduction in coordination. Inhibitions will be lowered causing one to potentially engage in risky behaviors.

0.08%: This is the current legal limit in ND that it is considered illegal and unsafe to drive.

0.10%: At this level, reaction time and control will be reduced, speech slurred, thinking and reasoning are slower and coordination of arms and legs is poor.

0.15%: This is a very high level with much less control over balance and voluntary muscles so walking and talking are difficult. May fall and vomiting begins.

0.20%: Stupor, confusion, feeling dazed and disorientation are common. Standing and walking may require help and sensations of pain decrease. Blackouts begin at this level as well as choking/aspiration on vomit.

0.30%: At this level, you may be unconscious and potential for death increases along with loss of understanding, increased heart rate, loss of bladder control.

0.40% and over: This level may put you in a coma or cause sudden death because your heart or breathing will suddenly stop.

Race

Select all

American Indian

Other

White

Sex

Select all

Female

Male

Age Group

Select all

10-19

20-29



30-39

40-49


50-59

60-69


70+



Alcohol Involvement by Year



Toxicology Present in Suicide Deaths



Race

Select all

American Indian

Other

White

Sex

Select all

Female

Male

Age Group

Select all

10-19

20-29



30-39

40-49

50-59

60-69

70+



Race

Select all

American Indian

Other

White

Sex

Select all

Female

Male

Age Group

Select all

10-19

20-29

30-39

40-49

50-59

60-69


70+




North Dakota ESSENCE Data



The Electronic Surveillance System for the Early Notification of Community-Based Epidemics (ESSENCE), captures syndromic surveillance data from hospitals in the State. This includes data from all acute care and critical access hospitals. This data consists of emergency department, urgent care and walk-in-clinic visit information. The purpose is to capture and analyze health-indicator data to identify abnormal health conditions, events, and enable early detection of outbreaks.


The data provided on the "Attempts" and "Ideation" pages, was obtained from the ND ESSENCE. Please note the caveats of this data:

1. These numbers represent a syndrome definition that utilizes both ICD-10 codes and chief complaint which looks for key words. These **should not** be considered a true "number of cases." Syndromes may also contain "noise" meaning that the syndrome data may count actual non-related events.
2. NOT every hospital submits both ICD and chief complaint so some visits may be missing.
3. Some hospitals only submit data on ND residents. Transient populations may not be included; therefore, underestimating the impact.
4. Increase in number may be due to actual increases or it may be due to increase in number of facilities participating.













Suicide Ideation by Sex



North Dakota Suicide Ideation



Suicide Ideation Rate per 10,000 by County Where Patient Lives





1000



1000

